

Estimation of steric sea level variations from combined GRACE and Jason-1 data

A. Lombard ^{a,*}, D. Garcia ^b, G. Ramillien ^a, A. Cazenave ^a, R. Biancale ^c, J.M. Lemoine ^c,
F. Flechtner ^d, R. Schmidt ^d, M. Ishii ^e

^a LEGOS-CNES, Observatoire Midi-Pyrénées, Toulouse, France

^b Space Geodesy Laboratory, Applied Mathematics Department, EPS, University of Alicante, Spain

^c Groupe de Géodésie Spatiale, CNES, Toulouse, France

^d GeoForschungsZentrum Potsdam, Potsdam, Germany

^e Frontier Research Center for Global Change, Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan

Received 3 August 2006; received in revised form 13 November 2006; accepted 17 November 2006

Available online 10 January 2007

Editor: C.P. Jaupart

Abstract

We estimate the mean steric sea level variations over the 60°S–60°N oceanic domain for the recent period (from August 2002 to April 2006), by combining sea level data from Jason-1 altimetry with time-variable gravity data from GRACE. The observed global mean sea level change from satellite altimetry results in total from steric plus ocean mass change. As GRACE measurements averaged over the ocean represents the ocean mass change component only, the difference between GRACE and altimetry observations provides an estimate of the mean steric sea level. Two different sets of GRACE geoid solutions (the GRGS EIGEN-GL04 and the GFZ EIGEN-GRACE04S products) have been used. Each GRACE data set ranges over approximately 3 yr or more (August 2002–April 2006 for the GRGS geoids and February 2003–February 2006 for the GFZ geoids).

We first focus on the seasonal variations. The two GRACE data sets agree very well in terms of mean annual mass variation, both in amplitude (approximately 7.3 mm equivalent sea level) and phase (maximum on October 14). For both time spans, Jason-1 sea level minus GRACE ocean mass has an annual amplitude of approximately 5.8 mm with a maximum on March 11. The latter signal compares well with the steric annual sea level estimated from the World Ocean 2004 climatology and the Ishii et al. [M. Ishii, M. Kimoto, K. Sakamoto, and S.I. Iwasaki, Steric sea level changes estimated from historical ocean subsurface temperature and salinity analyses, *Journal of Oceanography*, 62 (2) (2006) 155–170.] ocean temperature data. We also examine the interannual fluctuations of the Jason-1 minus GRACE sea level. The two resulting steric sea level time series (based on the two GRACE data sets) agree well. The inferred steric sea level curve exhibits an increasing trend during the last 3.5 yr (August 2002–April 2006), of the same order of magnitude as the 1993–2003 steric sea level trend computed with in situ hydrographic data. However, over the last 3.5 yr, we note a strong discrepancy between altimetry minus GRACE and in situ-based steric sea level trend, the latter exhibiting a negative slope. The cause for such a discrepancy is yet unknown but may be related to inadequate sampling of in situ ocean temperature measurements. © 2006 Published by Elsevier B.V.

Keywords: Ocean mass; Thermal expansion; Sea level; Climate change; Space gravimetry; Satellite altimetry

* Corresponding author. LEGOS-CNES, OMP, 18 Avenue E. Belin, 31401 Toulouse cedex 9, France. Tel.: +33 5 61 33 30 52; fax: +33 5 61 25 32 05.
E-mail address: Alix.Lombard@notos.cst.cnes.fr (A. Lombard).