

Moho depth variation in southern California from teleseismic receiver functions

Lupei Zhu

Southern California Earthquake Center, University of Southern California, Los Angeles

Hiroo Kanamori

Seismological Laboratory, California Institute of Technology, Pasadena

Abstract. The number of broadband three-component seismic stations in southern California has more than tripled recently. In this study we use the teleseismic receiver function technique to determine the crustal thicknesses and V_p/V_s ratios for these stations and map out the lateral variation of Moho depth under southern California. It is shown that a receiver function can provide a very good “point” measurement of crustal thickness under a broadband station and is not sensitive to crustal P velocity. However, the crustal thickness estimated only from the delay time of the Moho P -to- S converted phase trades off strongly with the crustal V_p/V_s ratio. The ambiguity can be reduced significantly by incorporating the later multiple converted phases, namely, the $PpPs$ and $PpSs+PsPs$. We propose a stacking algorithm which sums the amplitudes of receiver function at the predicted arrival times of these phases by different crustal thicknesses H and V_p/V_s ratios. This transforms the time domain receiver functions directly into the H - V_p/V_s domain without need to identify these phases and to pick their arrival times. The best estimations of crustal thickness and V_p/V_s ratio are found when the three phases are stacked coherently. By stacking receiver functions from different distances and directions, effects of lateral structural variation are suppressed, and an average crustal model is obtained. Applying this technique to 84 digital broadband stations in southern California reveals that the Moho depth is 29 km on average and varies from 21 to 37 km. Deeper Mohos are found under the eastern Transverse Range, the Peninsular Range, and the Sierra Nevada Range. The central Transverse Range, however, does not have a crustal root. Thin crusts exist in the Inner California Borderland (21–22 km) and the Salton Trough (22 km). The Moho is relatively flat at the average depth in the western and central Mojave Desert and becomes shallower to the east under the Eastern California Shear Zone (ECSZ). Southern California crust has an average V_p/V_s ratio of 1.78, with higher ratios of 1.8 to 1.85 in the mountain ranges with Mesozoic basement and lower ratios in the Mojave Block except for the ECSZ, where the ratio increases.

1. Introduction

The Mohorovicic discontinuity (Moho), which separates Earth’s crust from the underlying mantle, represents a major change in seismic velocities, chemical compositions, and rheology. The depth of Moho is an important parameter to characterize the overall structure of a crust and can often be related to geology and tectonic evolution of the region. Its lateral variation has strong influence on seismic wave propagation in the

crustal waveguide and controls the strong shaking from damaging earthquakes in certain distance ranges.

In southern California, tremendous efforts have been made to determine the Moho depth with a variety of geophysical methods. These include an early work of refraction studies using quarry blasts as energy sources [Kanamori and Hadley, 1975], seismic vertical reflection experiments in the Mojave Desert and vicinity [Cheadle et al., 1986; Li et al., 1992; Malin et al., 1995], tomographic imaging of Moho depth variation using Pn travel times from local earthquakes [Hearn, 1984; Hearn and Clayton, 1986; Sung and Jackson, 1992; Magistrale et al., 1992], teleseismic receiver function studies [Langston, 1989; Ammon and Zandt, 1993; Zhu and

Copyright 2000 by the American Geophysical Union.

Paper number 1999JB900322.
0148-0227/00/1999JB900322\$09.00