

Modelling of crustal anomalies of Lanzarote (Canary Islands) in light of gravity data

A. G. Camacho, F. G. Montesinos, R. Vieira and J. Arnosó

Instituto de Astronomía y Geodesia. (CSIC-UCM), Facultad de Ciencias Matemáticas, Ciudad Universitaria, 28040 Madrid, Spain.
E-mails: camacho@eucomax.sim.ucm.es; fuen@mat.ucm.es; vieira@mat.ucm.es; arnosó@mat.ucm.es

Accepted 2001 June 25. Received 2001 April 9; in original form 2000 July 20

SUMMARY

The application of a gravity inversion method enables us to obtain a 3-D density contrast model of the upper crustal anomalies of the volcanic island of Lanzarote (Canary Islands). For this, we use a network of 296 gravity stations distributed over the whole island, and a digital terrain model of about 45 000 terrestrial and oceanic data to determine the corresponding terrain correction. A density value of 2480 kg m^{-3} is chosen for this correction by means of a new approach. The resulting Bouguer anomaly is analysed by means of a least-squares prediction which gives us a mean level of uncorrelated observational noise of about 1.2 mgal. This anomaly is considered in order to obtain independent information about the inner anomalous mass density distribution by means of a 3-D gravity inversion based on a systematic exploration on a prismatic partition of the subsoil volume, and adopting *a priori* values of the density contrast (positive and negative) to determine the geometry of the anomalous bodies. The problem of non-uniqueness of the solution is avoided by using a minimization mix condition on the weighted residuals and the weighted whole anomalous mass. The structural solution is finally presented by means of horizontal sections and vertical profiles.

A main intrusive body is located under the central-eastern area and could correspond to a dilated volcanic activity of shield formation. It shows a prismatic form of more than 15 km depth, subducted with only the ridges remaining as horst blocks. Moreover, the SW and NE extreme areas of the island show smaller and shallower positive bodies, interpreted as less-developed magmatic intrusions. Conversely, several density lows offer interesting shallow alignments, 45°N (ENE–WSW) and 125°N (WNW–ESE), which could be associated with a fracture system corresponding to structural stress, and also correlate with historic eruptions, such as, for instance, the Timanfaya eruption. The monitoring of several geophysical parameters at two underground geodynamic stations, in the NE zone of the island and Timanfaya, shows characteristic differences between the two zones which confirm crustal anomalies in the second station.

Key words: gravity anomalies, inverse problem, Lanzarote island, volcanic structure.

INTRODUCTION

The Canary Islands (Fig. 1) are an old volcanic feature sited on top of Jurassic oceanic crust, located at the edge of the West African Continental Margin. Their origin is still under debate, and authors have proposed different types of genetic models (e.g. Wilson 1973; Schmincke 1982; Holik *et al.* 1991; Anguita & Hernan 1975; Marinoni & Pasquarè 1994) for the archipelago, where some age determinations (Abdel-Monem *et al.* 1971) indicate a general E–W age progression. The diverse models apply, for instance, hotspot theory (Wilson 1973), a connection

to the Alpine orogeny, which reached its maximum activity in this zone during the Miocene (Anguita & Hernán 1975), and the coexistence of a hotspot and a constraining complex regional structural framework (Schmincke 1982). Several geophysical research projects have been conducted in the archipelago (e.g. Bosshard & MacFarlane 1970; Banda *et al.* 1981; Canales & Dañobeitia 1998; Dañobeitia *et al.* 1994; Ranero *et al.* 1997), pointing out clear structural differences among the islands. Among these differences, from recent seismic studies it seems reasonable that the oceanic crust west of El Hierro islands is very shallow, Ranero *et al.* (1997) deduced a crustal thickness




Figure 1. Geographical location of Lanzarote island (29°N , 13.6°W ; 635000,3220000 UTM) at the northeastern extreme of the Canary archipelago (L: Lanzarote, A: Africa). Major volcanic periods in Lanzarote are (Marinoni 1991): (1) the Tableland (series I); (2) the Quaternary volcanism (series II, III); and (3) recent volcanism (series IV). Localities mentioned in the text: F, Famara; M, Mozaga; AJ, Los Ajaches; CV, Cueva de los Verdes; T, Timanfaya; SB, San Bartolomé; PC, Peñas del Chache; AF, Atalaya de Femés; CC, Caldera del Corazonzillo.

of between 6 and 9 km maximum, and beneath the continental shelf there are reported thicknesses of around 20–25 km (Hinz *et al.* 1982; Weigel *et al.* 1982).

The island of Lanzarote is located at the northeastern extreme of the archipelago, in clear alignment with the island of Fuerteventura (Fig. 1). The two islands constitute the emergent part of the East Canary Ridge, a NNE–SSW linear volcanic structure located on atypical oceanic crust, at least 11 km thick (Banda *et al.* 1981), between the continental rise and the Canary basin. The East Canary Ridge consists of several uplifted blocks of oceanic basement mantled by a thick pile of volcanic rocks (Grunau *et al.* 1975), probably overlying an intrusive complex. A sediment cover more than 10 km thick is located in the continental rise east of the ridge. Lanzarote is described by a shallow basement, probably about 4–5 km thick as deduced from seismic profiles (Banda *et al.* 1981), formed by a group of sedimentary rocks (quartzite and shales), plutonic rocks (basic and ultrabasic), and subvolcanic rocks (basaltic and trachytic dikes) with an abundance of xenoliths of quartzite and sandstone emitted by its volcanoes (Araña & Carracedo 1978). The lavas are basaltic, and very limited outcrops of massive thrahytes exist in the oldest parts of the island (NW and SE). A general geological study of Lanzarote can be found in Fuster *et al.* (1968) and Marinoni (1991). Three major volcanic periods are recognized (see Fig. 1): (1) the Tableland (series I) basalts dated at 6–12 Ma (Miocene–Pliocene); (2) the Quaternary volcanism (series II, III) dated at 1 Ma and separated from the former by an erosional interval; and (3) recent volcanism (series IV). Many Quaternary eruptions and several historic

ones have been identified. The Lanzarote eruption between 1730 and 1736 was one of the Earth's biggest historical eruptions. The structural evolution of this volcanic island may result from a complex interaction of magmatism with the regional stress field and the local stress field generated during the growth of the island itself.

According to Marinoni & Pascaré (1994), Lanzarote can be divided into three principal morphological blocks (Figs 1 and 2). The northernmost part is elongated in a NNE–SSW direction and consists mainly of the Famara massifs. The central part of the island has a characteristic flat truncated conic shape (Mozaga area). The southern block includes the Los Ajaches massifs. These remaining edifices correspond to the dilated eruptive episode of the shield stage (Miocene–Pliocene) with the first subaerial lava flows (Araña & Carracedo 1978). However, large earth movements may have disrupted this central structure until now, and only the ridges remain as horst blocks, the central cone having subsided between them and having been covered by younger volcanic eruptions (MacFarlane & Ridley 1969). Between the pre-erosional and the post-erosional stage, the Tableland succession was deeply eroded and probably tilted about 10° generally to the ESE (Marinoni & Pascaré 1994). After a long period of rest, volcanic activity was renewed (post-erosional stage during the Quaternary) and lasted almost until the present time; the resulting materials and volcanic edifices cover the major part of the island and have been well preserved.

So, the present structural architecture is the result of a complex magmatic and tectonic evolution characterized by variation in the stress field from the Miocene to the present.




Figure 2. Terrain model of the islands of Lanzarote and La Graciosa. Orthographic SW–NE view. Highest altitudes are about 600 m at Peñas del Chache (NE Lanzarote). In this figure two ancient and eroded edifices, Los Ajaches in the south and Famara in the north, are observed. One third central massif appears more eroded and subducted with respect to the former. In this central zone many well-preserved volcanic cones can be observed.

According to Armenti *et al.* (1989), the present structure is largely controlled by post-Miocene displacements which divide the island into five NW–SE elongated blocks, represented by the three structural highs (Famara, Mozaga and Los Ajaches) and two grabens. These blocks were subsequently dissected by NNE–SSW and NE–SW faults. The greatest principal stress shows an ENE–WSW direction, which argues for a transpressive regime. The NE–SW-striking faults were generated by extensional tectonics with a minor transurrence component. This resulted in a larger down-throwing of the northwestern sector relative to the southeastern one. A third narrow and elongated central graben, marked by the Timanfaya volcanic alignment, could be delineated between these two sectors.

Most of the geophysical works about the Canary Islands correspond to extended areas and marine data. Here we look for a local study corresponding to the upper crust of Lanzarote. To obtain a better understanding of the inner structure and evolution of Lanzarote, several geophysical and geodynamical studies were carried out on the island by the Instituto de Astronomía y Geodesia (IAG, Madrid). Two underground geodynamic stations monitoring several geophysical parameters were set up (Vieira *et al.* 1991; Arnosó *et al.* 2000). The station Cueva de los Verdes is located inside a lava tunnel of the La Corona volcano (Fig. 1) at the NE end of the island. The second station is located above the geothermal anomaly area in the Timanfaya National Park (Fig. 1). Shukowsky & Mantovani (1999) suggest that lateral variations in the scales of tectonic units could be inferred from tidal records. Continuous gravity tide observations have been carried out in these two laboratory locations for several years, and have been used to obtain the

respective local tidal gravity models (Table 1), which help in studying the anomalous crustal structure of the island in these two zones (Arnosó *et al.* 2000).

The IAG has also carried out several gravity studies on the island (e.g. Sevilla & Parra 1975). In this sense, gravity modelling plays an important role in studies of volcanic structures. Gravimetric methods have been employed by several authors (Malengreau *et al.* 1999; Rousset *et al.* 1989; Loddo *et al.* 1989; Camacho *et al.* 1997; Rymer & Brown 1986), and prove to be useful in the study of the origin, structure and activity of volcanic areas. The work of MacFarlane & Ridley (1969) corresponds to a smaller land gravity data set for Lanzarote. Gravity investigations do have, however, certain well-known limitations, the two most prominent perhaps being the deep resolution and the non-uniqueness of the solution. These disadvantages can be reduced by means of appropriate tools (geological information or statistical techniques).

Table 1. Observed amplitudes A and local phases α for the main tidal waves O_1 and M_2 at stations Cueva de los Verdes (coordinates: $29^\circ.16$ N, $13^\circ.44$ W, 37 m) and Timanfaya National Park (coordinates: $28^\circ.99$ N, $13^\circ.75$ W, 381 m). Amplitudes are given in μgal (10^{-8} ms^{-2}) and phases in degrees.

	Cueva de los Verdes		Timanfaya National Park	
	A	α	A	α
O_1	30.57 ± 0.02	-1.60 ± 0.04	29.74 ± 0.05	-1.45 ± 0.10
M_2	58.24 ± 0.02	2.27 ± 0.02	56.66 ± 0.03	2.07 ± 0.03

In 1988 a detailed gravity survey was carried out on Lanzarote island, the analysis of the data and preliminary structural model being presented in successive papers (Camacho & Vieira 1991; Camacho *et al.* 1992). The aim of the present paper is to obtain an improved interpretation of this gravity data by means of a 3-D inversion approach, to determine the geometry of the anomalous bodies corresponding to prescribed densities. This inverse method involves the systematic exploration of the model possibilities, admitting simultaneously positive and negative contrast (Camacho *et al.* 2000).

GRAVIMETRIC AND TOPOGRAPHIC DATA

The gravity data consist of 296 stations covering the islands of Lanzarote and La Graciosa with a nearly homogeneous distribution and a data separation of 1.5 km (Fig. 3). A LaCoste-Romberg gravimeter with digital electronic reading was used in the observation work, and a total of 437 observations were registered.

First, the gravity data were corrected taking into account the tidal model calculated using the records of the tidal gravity stations on the island (Arnosó 1996). Then, using the redundant observations, a global least-squares adjustment was made, taking as unknowns linear drifts, possible record jumps (as a result of mechanic or thermal shocks), and the (incremental) gravity values for each station. This adjustment produced residues with a standard deviation of 0.136 mgal ($1 \text{ mgal} = 10^{-3} \text{ m s}^{-2}$). Finally, the adjusted gravity data were referred to IGNS71. The station coordinates were determined by locating the points on detailed 1:5000 charts.

To calculate the further terrain correction of the gravity data, a digital terrain model was obtained for Lanzarote, La Graciosa and surrounding oceanic areas by means of a dense regular digitalization of 1:25 000 topographic charts. Fig. 2 shows an orthographic view of the resulting terrain model (45 000 data points). The highest altitudes are about

600 m. The figure shows two characteristic terrain elements. First, there are the two extreme areas of strongest relief: the Massif of Famara (northern) and the Massif of Los Ajaches (southern). These remaining edifices correspond to a dilated eruptive episode and posterior processes of large earth movements, subsidence, volcanic eruptions and an erosional stage. Second, there are the volcanic cones in the central area of Lanzarote and in La Graciosa. These preserved edifices, corresponding to the post-erosional stage, follow characteristic alignments.

Because of the proximity of Fuerteventura island, its contribution to the terrain correction was recalculated taking into account the terrain model of this island presented by Montesinos (1999).

GRAVITY ANOMALY

Using the 1980 normal gravity formula, and determining the terrain correction (extended to 45 km, according to the regularity of the effect beyond this distance), the refined Bouguer anomaly Δg was determined (Fig. 3). In choosing the density value for the terrain corrections we propose a new approach.

Several methods to calculate the terrain density, such as that of Nettleton (1939) which minimizes the correlation between the Bouguer anomaly and the topography, are efficient if the density of the formations is not related to the subsoil anomalous structure. In volcanic terrain, however, this correlation is generally predominant. The highest areas, for example, often correspond to the eruptive centres, which involve density anomalies.

An inadequate terrain density would assign an excess of topographical anomalous masses as sources of the anomaly field, producing some fictitious local gravity anomalies. Starting from the gravity data themselves, we look for a suitable terrain density that minimizes the gravity anomalies of topographic origin, keeping the anomalies due to the deep bodies. A tool to detect the presence of gravity anomalies of topographical




Figure 3. Distribution of gravity stations and map of the Bouguer gravimetric anomaly. The value used for the density of terrain correction is 2480 kg m^{-3} . Contour interval are 5 mgal. UTM coordinates in metres.

origin can be obtained by means of the anomalous masses model resulting from the inversion process. In fact, an inadequate terrain density will produce a fictitious increase of the local anomalies. It will cause an excess of (shallow) anomalous masses as sources of the anomaly field. Therefore, the proposed criterion is to choose the density that produces a minimum value of the whole anomalous mass of the model that results from applying the inversion approach.

To apply this criterion, however, we must take into account that, in the inversion process (see next section), the whole resulting magnitude of the anomalous mass also depends on the fit level (mean resulting residues) selected for the gravity anomalies (which, in turn, is regulated by means of a factor in the inversion process). Thus, to reach this desirable minimum of mass, we test the gravity inversion results corresponding to several terrain densities and to several possible fit levels (mean residual level).

This method was applied to the gravity anomalies of Lanzarote, resulting in a mean value of 2480 kg m^{-3} for the terrain mass density. Fig. 4 shows the results of various inversion tests (according to the inversion method detailed below) corresponding to several possible terrain correction densities and to different allowed mean residues. The chosen value is an intermediate density somewhat greater than 2300 kg m^{-3} , suggested by MacFarlane & Ridley (1969) as a realistic value for the upper layer of the crustal structure from seismic refraction data, but smaller than the 2560 kg m^{-3} that can be obtained by Nettleton's method (null correlation altitude-anomaly).

With the adopted terrain density, the terrain corrections range between 2 and 12 mgal with a mean value of 3.3 mgal, and the resulting Bouguer anomaly ranges between 152 and 223 mgal (Fig. 3).

A study of the Bouguer anomaly by means of covariance analysis of the data and a further least-squares prediction (e.g. Montesinos *et al.* 1999) enable us to filter the data and obtain a value 1.2 mgal as the standard deviation of the non-correlated noise (Fig. 5). The value is conditioned not only by the accuracy of the (gravimetric and topographic) data, but also by the distance (about 1.5 km) between stations and the horizontal




Figure 4. Relation between resulting whole anomalous mass ($\times 10^9 \text{ kg}$) and various allowed mean residues (standard deviation, mgal) for several inversion tests corresponding to several possible terrain correction densities. The value of 2480 kg m^{-3} provides the minimum whole anomalous mass and is therefore selected for the terrain correction.




Figure 5. Histogram of the residues of the filtering process. This non-correlated noise of the data follows a normal distribution of mean 0 mgal and standard deviation 1.19 mgal.

gravity gradients (about 3 mgal km^{-1}) present in the area. This method enables us to obtain a suitable covariance matrix, corresponding to the inaccuracies for the gravity data, that is useful in the following inversion process.

GRAVITY INVERSION METHOD

While limited by the validity of the adopted hypothesis, the gravity inversion methods that seek to determine the geometrical properties of anomalous bodies with fixed density contrast correspond to a non-linear environment and offer interesting results. Usually these methods (linearized approaches, iterative gradient methods, etc.) start from good initial solutions and improve iteratively (e.g. Barbosa *et al.* 1997; Enmark 1981). General complications come from adopting a 3-D environment and when positive and negative anomalous contrasts are simultaneously accepted. In addition, the absence of a good initial model combined with data inaccuracies and uncertainties concerning regional trend introduce more ambiguity into the inverse problem. For the fully non-linear treatment, the methods of exploration of the space model are often the best option (Tarantola 1987). This exploration process can be conducted randomly (Silva & Hohmann 1983) or by means of a systematic approach, according to the size of the model space to explore. Here we consider an inversion method, a modification of the one described in Camacho *et al.* (2000), which can be included in this last group of systematic exploration.

For this non-linear method, the usual strong hypothesis is adopted: the subsurface anomalous structure is characterized by prescribed mass density contrasts. Moreover, positive and

negative density contrasts are simultaneously accepted. Therefore, the problem consists of determining the geometry of the anomalous volumes corresponding to those prescribed density contrasts, and responsible for the observed gravity anomaly. To describe that geometry, the subsurface volume is divided into a fixed discrete 3-D partition of parallelepiped cells, and then the anomalous volumes are constructed and described as aggregations of elements filled with the prescribed density contrasts.

The main problem for the applicability of the exploratory methods is the large size of the space model to be explored. In our case, a direct application (test across every possible geometrical configurations) of the systematic exploratory search would be very tedious. This is solved by using a ‘growth’ process or expansion approach to construct the anomalous bodies. In this case, the exploration of every model possibility is substituted by exploration of several possibilities of growth (cell by cell) for each step of the expansion of the anomalous bodies. This is a more reasonable goal. Thus, step by step, the prismatic cells, each with prescribed densities, are systematically tested, and then the best options are adopted and added to the growth approach for the anomalous bodies construction. Simultaneously, a regional trend $g_{\text{reg}}(x, y)$ of the gravity data can be adjusted.

For the k th step of the growth process, the residues v_i , $i = 1, \dots, N$, for the N gravity stations can be calculated as

$$v_i = \Delta g_i - f \sum_{j \in J_k} A_{ij} \Delta \rho_j + g_{\text{reg}}(x_i, y_i), \quad i = 1, \dots, N, \quad (1)$$

where Δg_i is the observed anomaly; A_{ij} is the vertical attraction of the j th cell on the i th gravity station for unit density; $\Delta \rho_j$ is the density contrast value assigned to the j th cell (among the possible prescribed values); J_k is the set of indexes corresponding to the cells filled until this k th step and then constituting the growing anomalous body; x_i, y_i are the station coordinates; and $f \geq 1$ is an adjusted scale factor for the fit.

If only a minimization criterion for the residuals \mathbf{v} is used, the acceptance of positive and negative values for the prescribed density contrasts and the inclusion of the trend unknowns give a non-uniqueness problem. To remedy this, an additional condition of minimization of the model variation can be adopted. Thus, the solution is obtained by a mixed condition formed by the gravity fit with the l_2 norm and the whole anomalous mass quantity, using the parameter λ for a suitable balance:

$$\mathbf{v}^T \mathbf{Q}_D^{-1} \mathbf{v} + \lambda f^2 \mathbf{m}^T \mathbf{Q}_M^{-1} \mathbf{m} = \min., \quad (2)$$

where $\mathbf{v} = (v_1, \dots, v_N)^T$ are the gravity residues given by (1) for the N stations, $\mathbf{m} = (\Delta \rho_1, \dots, \Delta \rho_{NB})$ are the anomalous densities for the NB cells of the subsoil partition, λ is a positive factor, empirically fixed, for balance between model fitness and model smoothness, and $f \geq 1$ is the adjustable scale factor which keeps the model gravity nearly proportional to the observed value. \mathbf{Q}_D is the covariance matrix (usually a diagonal matrix) corresponding to the estimated (Gaussian) inaccuracies of the gravity data, and \mathbf{Q}_M is a covariance matrix corresponding to the supposed determinability of the model parameters \mathbf{m} . We take, as covariance matrix \mathbf{Q}_M , a diagonal normalizing matrix of non-null elements the same as the diagonal elements of $\mathbf{A}^T \mathbf{Q}_D^{-1} \mathbf{A}$. The λ parameter governs the application of the minimization conditions regarding the balance between total anomalous mass and residual values (Fig. 4). For low λ -values

a better fit is obtained, but the anomalous mass may increase excessively and adopt too deep a position. For high λ -values the adjusted model can be too poor and shallow. The λ -value is chosen so that the standard deviation of the inversion residues is similar to the data noise level (1.2 mgal for the data of Lanzarote) determined by the previous covariance analysis.

For each step of the anomalous growth, a systematic exploration of the expansion possibilities is tested with respect to the value given by (2), determining also the corresponding f -value and the parameters of a regional component adjusted as a linear trend. This expansion process stops when the scale factor f is close to one. Finally, the solution appears as a 3-D distribution of prismatic cells that have been assigned some of the prescribed contrast densities. Moreover, a regional trend is also obtained, complementing this anomalous mass distribution.

This inversion method, detailed in Camacho *et al.* (2000), requires, as usual for many non-linear methods, a suitable choice of the *a priori* density contrasts (positive and negative in our case). A too-high density contrast would fit the main anomaly component but would produce a rather ‘skeleton’ model, perhaps unable to represent very slight anomalous structures. A too-low density contrast would fit the light structures, but would produce a rather ‘inflated’ model, perhaps unable (taking into account the volume limitations) to fit the main components, and giving rise to fictitious peripheral bodies. To remedy this, and if geological knowledge of the true local density contrasts is not available, we propose an improvement of our inversion method by means of a theoretical approach.

Instead of invariable density contrasts along the whole model formation, we propose now to adopt changing contrasts. Starting from fixed maximum values (let us name them R_P positive and R_N negative), the prescribed density contrasts evolve during the growth process according to a simple law. By means of empirical tests, we have selected for each growth step the density contrasts $\Delta \rho_P$, and $\Delta \rho_N$, positive and negative, given by

$$\Delta \rho_P = R_P e^{-f^\tau}, \quad \Delta \rho_N = R_N e^{-f^\tau}, \quad (3)$$

where $f \geq 1$ is the scale factor corresponding to the step of the fit process, and τ is a fixed factor corresponding to the desired variability of the density values. A high τ -value will produce an anomalous model with contrast values mostly close to the maximum values (and thus with a sharp geometry). A low τ -value will produce a model with more variable or decreasing contrasts (and thus with a more diluted geometry). In this form, the high density contrasts (close to the maximum values R_P and R_N) will be employed to fit the main anomaly components, while smaller density contrasts fit the smaller and local anomalies. The resulting anomalous model is richer (and perhaps somewhat more diffuse).

We must keep in mind that the gravity inversion is nearly insensible to horizontal mass stratification, and therefore the obtained model must be considered independent of any additional stratification. In spite of these limitations, this method provides interesting gravimetric information. The method has the following advantages:

- (1) it adopts a 3-D environment;
- (2) non-gridded, non-planar and inaccurate data are accepted;
- (3) a simple regional trend can be simultaneously determined;
- (4) adjustment of an indefinite number of anomalous bodies is possible;

- (5) non-regular subsoil partition can be considered (e.g. with deeper blocks bigger than shallow blocks);
- (6) if previous evaluated models exist, they can be incorporated;
- (7) a variable density contrast can be adopted in the inversion process; and, above all
- (8) positive and negative density contrasts are simultaneously accepted (see Camacho *et al.* 2001 for software development).

ADJUSTED 3-D MODEL OF ANOMALOUS DENSITY

Taking into account that the distance between contiguous stations is about 1500 m and that the mean diameter of the survey is about 30 000 m (and the computer limitations), we adopted a 3-D partition of the local subsoil into 16 800 rectangular prisms, with sides ranging from 700 m for the shallow elements (1000 m depth) to 1300 m for the deepest blocks (at 18 000 m depth). According to the suggested method, we selected, for the whole subsoil volume, the possible density contrasts by means of the extreme values, $R_N = -440 \text{ kg m}^{-3}$ and $R_P = +440 \text{ kg m}^{-3}$, and an exponential variation law (with $\tau=4$) of the contrasts. These values have been selected empirically to obtain connected and well-developed volumes and to produce an inversion model with a negative mean density contrast of -346 kg cm^{-3} and a positive one of 341 kg cm^{-3} , which could correspond to the usual contrasts among structural elements of the Canarian volcanism (these values are close to the contrasts that occur when considering, for instance, density values of 2250 kg m^{-3} for sedimentary material, 2700 kg m^{-3} for crustal material, and 3100 kg m^{-3} for mantle-type material) (Bosshard & MacFarlane 1970). When different density contrasts are used, the resulting solutions are rather similar, although for higher contrasts, smaller volumes would be obtained.

Given this partition of the subsoil, and considering the above density contrasts, we applied our inversion method and obtained a model with the geometry of the anomalous bodies (for the adopted densities) and a simple regional component of the gravity anomaly field by means of a polynomial surface of degree one. This component reflects the general tendency of the anomaly and is strongly conditioned by the two main anomaly highs located in the east and northeast of the island.

The adjusted local model of density contrasts is shown by several horizontal sections and vertical profiles in Figs 6 and 7, from which some interesting features can be highlighted.

The main structural element in the adjusted model is a positive contrast mass ('A' in Fig. 8) which emerges close to the centre of the island, in the area of San Bartolomé (see Fig. 1). Its lateral size is about 13 km (Fig. 6), and, according to the adjusted model, its anomalous mass is about $582 \times 10^{12} \text{ kg}$. The top of this body, located in the shallower sections of the model, presents a double summit ('A1' and 'A2' in Fig. 9) with both summits elongated along 45°N and with a mutual strike of 125°N . Its bottom appears in the adjusted model with a depth greater than 15 km (Figs 6 and 7) (not far from the supposed depth of the Moho). The maximum depth of this body is, however, somewhat doubtful, because its determination interferes with the regional trend adjustment. While for the deep sections (under 7 km) this main body appears rather rounded in the model, for mean depths (1–6 km), where the gravity

survey is more sensitive, the model shows a clear geometry with a nearly square form according to directions 45°N (NNE–SSW) and 125°N (NNW–SSE). This important anomalous mass with its nearly prismatic shape can be interpreted as an intrusive body. According to MacFarlane & Ridley (1969), this body corresponds to a major igneous centre within the crust; it was important at least during the early subaerial growth of the island, and may have formed initially at the intersection of fundamental NNE- and SW-trending fault systems. However, while large earth movements and an erosional process may have disrupted this structure until now, only the ridges remain as horst blocks, the central cone having subsided between them and having been covered by younger volcanic eruptions. This hypothesis is in agreement with a possible interpretation of our model taking into account the irregular form (with negative and positive contrasts) in the shallower sections, associated with recent volcanism and the presence of the two structures A1 and A2. Marinoni & Pasquare (1994) relate this intrusive body to surface outcrops of the Shield phase. An alternative explanation of the very superficial low between A1 and A2 could be a simple erosion process.

A subsidiary intrusive body is located from 1 to 6 km depth close to the main body (body B in Fig. 8), with an anomalous mass of about $33 \times 10^{12} \text{ kg}$. This structure suggests a much smaller accumulation of magma in the crust, drained from the central reservoir (A in Fig. 8) along a SSW–NNE-trending fissure system indicated as the alignment of the two bodies. The system follows the WSW–ENE direction of the principal stress in the island (Armenti *et al.* 1989). The body is situated to the south of the Peñas del Chache, in the Macizo de Famara (see Figs 1 and 2), the highest point of the island and one of the oldest areas.

Two positive bodies are also noticeable: a peripheral intrusive body (C in Fig. 8), located SW of Lanzarote and elongated along 126°N ; and a third intrusive body (D in Fig. 8) located at the NE extreme of Lanzarote and elongated along 133°N . This last body is too peripheral for its limits to be determined, although the adjusted model suggests a magnitude smaller than that of the main body. The gravity data observed in La Graciosa Island do allow us, however, to define the geometry of this body on its SW side.

Body C lies west of Atalaya de Femés volcano (see Fig. 1), which was constructed in the post-erosional stage (Marinoni & Pasquare 1994). It extends between the shallower sections of the model and 7 km depth with an anomalous mass of $48 \times 10^{12} \text{ kg}$. A more superficial and smaller body appears aligned along 43°N (E in Fig. 8), close to several small outcrops of Series I (Coello *et al.* 1992).

The existence of these two less developed and shallower positive bodies, C and D, suggests that the accumulation of mantle-type material took place close to the surface. Between these not very massive SW and NE areas, the central body, A, suffered a further subsidence process during growth (Watts 1994), producing an extensive body with deep roots as is observed in the deeper sections. According to this interpretation, MacFarlane & Ridley (1969) note that the topography and geology of the island suggest that the central part has been down-faulted between the uplands in the north and south, forming a graben structure.

All these positive bodies present a nearly regular prismatic shape (within the intrinsic smoothness of shape for the deep zones in the inversion of potential fields), with borders along




Figure 6. Adjusted model of anomalous density contrast obtained by means of 3-D gravity inversion. Several horizontal sections, from -2000 to $-14\,000$ m depth, are shown.

45°N and 125°N . Moreover, the adjusted model shows others characteristic alignments. The main body, A, shows a like-alignment with the bodies B and D, with strike about 23°N . Meanwhile, the bodies in the south of the island (C and E) lie, with similar strike, along a line 15 km displaced to the west, and they could be connected by alignment with the crustal structures identified in the north and central area of the neighbouring island Fuerteventura by means of a similar gravimetric interpretation (Montesinos 1999). So, MacFarlane & Ridley (1969) also suggest lateral movements as inferred by

the NNE-trending scrap visible in the northern uplands that appears to have been displaced $15\text{--}20$ km westwards in the south of the island.

Two main kinds of low-density areas are shown in the adjusted model: extensive, peripheral and deep minimum structures that limit the E and SE borders of the island; and the internal minimum areas which form elongated trenches.

The first group, extrapolated in the model, and mostly on the west side of the island, may be associated with oceanic sediments coming from the nearby African margin that infill




Figure 7. Adjusted model of anomalous density contrast obtained by means of 3-D gravity inversion. Several vertical profiles across the main structures (according to the lines shown in Fig. 6d) are shown.

the moat as a result of the subsidence of the main intrusive body A. We observe (Figs 6 and 7) that these structures appear from the surface down to a depth of about 7 km. However, taking into account the extrapolated nature of these lows, reliable geometric properties cannot be deduced.

The second group is composed of thin elongated structures that run across the centre and western area of the island with depths ranging from 1 to 4 km. These trenches follow two main directions, 45°N(ENE–WSW) and 125°N(WNW–ESE), across the island (Fig. 9). The Timanfaya fissural eruption occurred (1730–1760) within this minimum-density area; its fractures and volcanic alignment followed two main directions, E–W and ENE–WSW, each one defined by smaller alignments of superposed vents (Marinoni & Pasquaré 1994). Connected with this trench pattern, we observe a main minimum of greater depth (Fig. 8) under the cone of Caldera del Corazoncillo (see Fig. 1), in the zone of Timanfaya, at the intersection of two alignments of minima according to the main directions described (Fig. 9).

Other peripheral trenches of low density with local directions are also identified in the model; for instance, with a direction of

nearly N–S in Famara, and a direction of nearly E–W in the northwestern part of the island (Figs 6 and 9). In La Graciosa, another small local minimum is identified in the eastern part of the island. We consider all these elongated low-density structures to be connected with fractured areas and recent volcanism.

The obtained model is in agreement with the hypothesis that the central area of Lanzarote is a subducted structure, crossed by a system of shallow fractures, indicating a crustal difference with respect to peripheral formations at Famara and Los Ajaches (e.g. Armenti *et al.* 1989). The NW zone of this central area, and especially Timanfaya, appears to be subjected to the main fractures deduced from the adjusted model. These crustal differences have also been observed by means of several techniques that highlight discrepancies between results obtained in Timanfaya and other results in the zone of Famara (Laboratory of Cueva de los Verdes). For instance, the gravity tide anomalies computed after correcting the ocean loading and attraction effects (Arnosó *et al.* 2000) show clear differences in magnitude at the two stations, being larger for M_2 and O_1 at Timanfaya than at Cueva de los Verdes station. Moreover, the phases of




Figure 8. Horizontal section of the anomalous density contrast model corresponding to a model mean depth of 4000 m. The fairly regular geometry of the positive density contrast bodies is highlighted. These structures are interpreted as a system of intrusive bodies with SWS–ENE trend. Subsidiary bodies thus appear as drained from the main sources (the central body A and the smaller structure B).

the residuals for O_1 and M_2 are more consistent with a body tide effect than with an ocean loading effect, and the sign of the O_1 and M_2 residuals at Timanfaya station are negative, which indicates a response of a porous or fractured local upper crust subject to the influence of tidal strain (Arnosó *et al.* 2001). Díez Gil *et al.* (1986), by means of thermal anomalies, suggest the

existence of a magmatic intrusion, of radius 200 ± 100 m, located at a depth of 4 ± 1 km, with a convective system in which the energy is transported by fractures which continue to depth according to electromagnetic studies. These results are compatible with air model and with the existence of bodies of negative density contrast such as those indicated.




Figure 9. Shallow horizontal section of the anomalous density contrast model corresponding to a depth of 2000 m. A pattern of alignments of negative density contrast zones, with main directions ENE–WSW and WNW–ESE, is identified and associated with fracture areas and recent volcanism.

Finally, Fig. 10 shows the distribution of residues of the inversion adjustment, which have a standard deviation of 1.15 mgal, in accordance with the data accuracy.

CONCLUSIONS

The application of a new version of the gravity inversion method proposed in Camacho *et al.* (2000) on the gravity data observed in Lanzarote and La Graciosa enables us to obtain a model of crustal anomalous mass for these islands. The gravity values considered have been analysed, resulting in a level of data accuracy of 1.2 mgal (non-correlated noise). To obtain the Bouguer anomaly we use a density for terrain correction of 2480 kg m^{-3} . This value is calculated by means of a new approach based on the whole anomalous mass resulting for the inversion model. The obtained Bouguer anomaly ranges between 152 and 223 mgal, and is used to apply the modified inversion method. The actual improvement of the method consists of permitting the variability of the density contrasts that appear in the model. The adjusted model for Lanzarote is constructed by limiting the possible density contrasts to between -440 and 440 kg m^{-3} , resulting in a geometrical configuration of high- and low-density areas. The results show a main intrusive body as a result of the accumulation of mantle-type material and further subsidence. Other smaller and shallow intrusive bodies are detected. These positive bodies can be related to the shield stage of the island formation. Their geometry is characterized by regular shapes with clear strikes (45°N and 125°N). This regularity suggests a tectonic component in the origin and evolution of these structures. The adjusted low-density bodies




Figure 10. Distribution of residues corresponding to the inversion process. A standard deviation of 1.15 mgal, close to the value obtained for the data noise, is presented.

comprise some peripheral structures and several linear trenches which we associate with a regular fracture system. Some of these crustal anomalies have been confirmed using results obtained in two geodynamic laboratories (in Timanfaya and Cueva de los Verdes). The gravity tide anomalies show a larger magnitude at Timanfaya than at Cueva de los Verdes station, and the results at Timanfaya station indicate the existence of fractured and porous upper crust.

ACKNOWLEDGMENTS

This work is part of the projects PB88-0022, PB91-0150 and AMB97-0706 conducted by the Instituto de Astronomía y Geodesia and developed in the Canary Islands. Additional support was obtained from the Cabildo Insular de Lanzarote. J. J. Dañobeitia provided helpful comments on the manuscript.

REFERENCES

- Abdel-Monem, A., Watkins, N.D. & Gats, P.W., 1971. Potassium argon ages, volcanic stratigraphy, and geomagnetic polarity history of the Canary islands: Lanzarote, Fuerteventura, Gran Canaria and La Gomera, *Am. J. Sci.*, **271**, 490–521.
- Anguita, F. & Hernán, F., 1975. A propagating fracture model versus a hot spot origin for the Canary Islands, *Earth planet. Sci. Lett.*, **27**, 11–19.
- Araña, V. & Carracedo, J.C., 1978. *Canarian Volcanoes. II Lanzarote y Fuerteventura*, Rueda, Madrid.
- Armenti, P., Marinoni, L. & Pasquare, G., 1989. Geological Map of the Island of Lanzarote, in *ESF Meeting on Canarian Volcanism*, pp. 198–200, European Science Foundation, Madrid.
- Arnosó, J., 1996. Modelización y evaluación de efectos indirectos sobre las mareas terrestres en el área de las Islas Canarias, *PhD thesis*, University Complutense of Madrid, Spain.
- Arnosó, J., Fernández, J., Vieira, R. & Ruymbeke, M., 2000. A preliminary discussion on tidal gravity anomalies and terrestrial heat flow in Lanzarote (Canary Islands), *Bull. Inf. Marées Terrestres*, **132**, 10 271–10 282.
- Arnosó, J., Fernández, J. & Vieira, R., 2001. Interpretation of tidal gravity anomalies in Lanzarote, Canary Islands, *J. Geodyn.*, **31**, 341–354.
- Banda, E., Dañobeitia, J.J., Suriñach, E. & Ansonge, J., 1981. Features of crustal structure under Canary Islands, *Earth planet. Sci. Lett.*, **55**, 11–24.
- Barbosa, V.C.F., Silva, J.B.C. & Medeiros, W.E., 1997. Gravity inversion of basement relief using approximate equality constraints on depths, *Geophysics*, **62**, 1745–1757.
- Bosshard, E. & MacFarlane, D.J., 1970. Crustal structure of the W Canary Islands from seismic refraction and gravity data, *J. geophys. Res.*, **75**, 4901–4918.
- Camacho, A.G. & Vieira, R., 1991. Gravimetric study of Lanzarote Island, in *Cahiers du Centre European de Geodynamique et de Sismologie. Conseil l'Europe*, **4**, 339–351.
- Camacho, A.G., Vieira, R., Montesinos, F.G. & Arnosó, J., 1992. Investigaciones gravimétricas sobre la estructura de la corteza volcánica en la isla de Lanzarote, in *Int. Conf. on Cartography-Geodesy, Maracaibo (Venezuela) II*, pp. 364–372, eds Sevilla, M.J., Henneberg, H. & Vieira, R., I.A.G. (CSIC-UCM), Madrid.
- Camacho, A.G., Montesinos, F.G. & Vieira, R., 1997. A three-dimensional gravity inversion applied to S. Miguel island (Azores), *J. geophys. Res.*, **102**, 7717–7730.
- Camacho, A.G., Montesinos, F.G. & Vieira, R., 2000. Gravity inversion by means of growing bodies, *Geophysics*, **65**, 95–101.
- Camacho, A.G., Montesinos, F.G. & Vieira, R., 2001. A 3-D gravity inversion tool based on exploration of model possibilities, *Comput. Geosci.*, in press.

- Canales, J.P. & Dañoibetia, J.J., 1998. The Canary Islands swell: a coherence analysis of bathymetry and gravity, *Geophys. J. Int.*, **132**, 479–488.
- Coello, J., Cantagrel, J.M., Hernán, F., Fúster, J.M., Ibarrola, E., Ancochea, E., Casquet, C., Jamond, C., Díaz de Téran, J.R. & Cendrero, A., 1992. Evolution of the eastern volcanic ridge of the Canary Islands based on new K-Ar data, *J. Volc. Geotherm. Res.*, **53**, 251–274.
- Dañoibetia, J.J., Canales, J.P. & Dehghani, G.A., 1994. An estimation of the elastic thickness of the lithosphere in the Canary Archipelago using admittance functions, *Geophys. Res. Lett.*, **21**, 2649–2652.
- Díez Gil, J.L., Juguero, J., Ortiz, R. & Araña, V., 1986. Termometrias y modelos matemáticos para el estudio de gradientes térmicos superficiales en Lanzarote (Islas Canarias), *Anal. Física B*, **82**, 91–101.
- Enmark, T., 1981. A versatile interactive computer program for computation and automatic optimization of gravity models, *Geoexploration*, **19**, 47–66.
- Fuster, J.M., Fernandez Santin, S. & Sagredo, J., 1968. *Geology and Volcanology of the Canary Islands*. Lanzarote, Instituto Lucas Mallada, CSIC, Madrid.
- Grunau, H.R., Lehner, P., Cleintuar, M.R., Allebach, P. & Bakker, G., 1975. New radiometric ages and seismic data from Fuerteventura (Canary Islands), Maio (Cape Verde Island) and Sao Tomé (Gulf of Guinea), in *Progress in Geodynamics*, pp. 90–118, eds Borradaile, G.J. et al., R. Soc. Neth. Acad. Arts, Sci., Amsterdam.
- Hinz, K., Dostmann, H. & Frisch, J., 1982. The continental margin of Morocco: seismic sequences, structural elements and geological development, in *Geology of the Northwest African Continental Margin*, pp. 34–59, eds von Rad, U., Hinz, K., Sarnthein, M. & Seibold, E., Springer, Berlin.
- Holik, J., Rabinowitz, P. & Austin, J.A., 1991. Effects of canary hotspots volcanism on structure of oceanic crust off Morocco, *J. geophys. Res.*, **96**, 12 039–12 067.
- Loddo, M., Patella, R., Quarto, G., Ruina, A., Tramacere, A. & Zito, G., 1989. Application of gravity and deep dipole geoelectrics in the volcanic area of Mt. Etna (Sicily), *J. Volc. Geotherm. Res.*, **39**, 17–39.
- MacFarlane, D.J. & Ridley, W.I., 1969. An interpretation of gravity data for Lanzarote, Canary Islands, *Earth planet. Sci. Lett.*, **6**, 431–436.
- Malengreau, B., Lenat, J.F. & Froger, J.L., 1999. Structure of Reunion Island (Indian Ocean) inferred from the interpretation of gravity anomalies, *J. Volc. Geotherm. Res.*, **88**, 131–146.
- Marinoni, L.B., 1991. Evoluzione geológica e strutturale dell'isola di Lanzarote (Islas Canarias), *PhD thesis*, University of Milan, Milan.
- Marinoni, L.B. & Pasquarè, G., 1994. Tectonic evolution of the emergent part of a volcanic ocean island: Lanzarote, Canary Islands, *Tectonophysics*, **239**, 111–135.
- Montesinos, F.G., 1999. Inversión gravimétrica 3D por técnicas de evolución. Aplicación a la isla de Fuerteventura, *PhD thesis*, University Complutense of Madrid, Spain.
- Montesinos, F.G., Camacho, A.G. & Vieira, R., 1999. Analysis of gravimetric anomalies in furnas caldera (Sao Miguel, Azores), *J. Volc. Geotherm. Res.*, **92**, 67–81.
- Nettleton, L.L., 1939. Determination of density for reduction of gravimeter observations, *Geophysics*, **4**, 176–183.
- Ranero, C.R., Banda, E. & Buhl, P., 1997. The crustal structure of the Canary basin: Accretion process at slow spreading centers, *J. geophys. Res.*, **102**, 10 185–10 201.
- Rousset, D., Lesquer, A., Bonneville, A. & Lenat, J.F., 1989. Complete gravity study of Piton de la Founaise volcano, Reunion Island, *J. Volc. Geotherm. Res.*, **36**, 37–52.
- Rymer, H. & Brown, G.C., 1986. Gravity fields and the interpretation of volcanic structures: Geological discrimination and temporal evolution, *J. Volc. Geotherm. Res.*, **27**, 229–254.
- Schmincke, H.U., 1982. Volcanic and chemical evolution of the Canary Islands, in *Geology of the Northwest African Continental Margin*, pp. 273–301, eds von Rad, U., Hinz, K., Sarnthein, M. & Seibold, E., Springer, Berlin.
- Sevilla, M. & Parra, R., 1975. Levantamiento gravimétrico de Lanzarote, *Rev. R. Ac. CC. Exactas, Físicas y Naturales*, **69**, 257–284.
- Shukowsky, W. & Mantovani, M.S.M., 1999. Spatial variability of tidal gravity anomalies and its correlation with the effective elastic thickness of the lithosphere, *Phys. Earth planet. Inter.*, **114**, 81–90.
- Silva, J.B.C. & Hohmann, G.W., 1983. Nonlinear magnetic inversion using a random search method, *Geophysics*, **46**, 1645–1658.
- Tarantola, A., 1987. *Inverse Problem Theory. Methods for Data Fitting and Model Parameter Estimation*, Elsevier, Amsterdam.
- Vieira, R., Van Ruymbeke, M., Fernández, J., Arnoso, J. & Toro, C., 1991. The Lanzarote underground laboratory, *Cahiers du Centre Européen Géodynamique de Séismologie*, **4**, 71–86.
- Watts, A.B., 1994. Crustal structure, gravity anomalies and flexure of the lithosphere in the vicinity of the Canary Islands, *Geophys. J. Int.*, **119**, 648–666.
- Weigel, W., Wissmann, G. & Goldflam, P., 1982. Deep seismic structure (Mauretania and Central Morocco), in *Geology of the Northwest African Continental Margin*, pp. 132–159, eds von Rad, U., Hinz, K., Sarnthein, M. & Seibold, E., Springer, Berlin.
- Wilson, J.T., 1973. Mantle plumes and plate motions, *Tectonophysics*, **19**, 149–164.