


Part IV: Depositional Environments

A depositional environment is a **geomorphic unit** in which deposition takes places. Thus, the study of depositional environment is essentially the study of geomorphology, i.e. **recognition of geomorphic units**. Geomorphic units are recognized by features preserved in ancient sediments.



沈積環境示意圖。陸棚水深200m以內，陸坡深200m至洋盆。注意陸坡坡度只有4-5度，本圖因有垂直放大，使陸坡坡度似達80度。(鄧屬予，1997)

Principles of environmental interpretation and classification

Facies Analysis

- 1. Definition:** A **facies** is a body of rock characterized by a particular combination of lithology, texture, suite of sedimentary structures, fossil content, colour, geometry, paleocurrent pattern, etc.
- 2. How is a facies formed?** A facies is produced by one or several processes operating in a depositional environment.
- 3. Description:** Facies are best referred to objectively in purely descriptive terms, using a few pertinent adjectives; examples could be cross-bedded, coarse sandstone facies or massive pebbly mudstone facies.
- 4. Facies association:** Groups of facies genetically related to one another and which have some environmental significance can be designate as a **facies association**. The facies comprising an association are generally deposited in the same broad environment, in which there are several different depositional processes operating.

Facies analysis is to identify geomorphic units in which ancient sediments accumulated.

Sedimentary Environment (Process Element)

Dynamic elements of the environment

Physical processes: wave and current activity; gravity processes; sea-level changes; tectonism and volcanism

Chemical processes: solution, precipitation, authigenesis

Biological processes: biochemical precipitation; biologic reworking of sediment; photosynthesis

Static elements of the environment

Geomorphology of the depositional site

Water depth

Water chemistry

Depositional materials (sediment supply)

Climate



Sedimentary Facies (Response Element)

Geometry of the deposit

Blanket, prism, shoestring, etc.

Primary sediment properties

Physical: bedding and contact relationships; sedimentary textures and structures; color; particle composition

Chemical: major-element and trace-element composition

Biological: fossil content (types and abundances of fossils)

Derived sediment properties

Porosity and permeability

Acoustical properties (sound transmissibility)

Electrical conductivity

Radioactivity


An example on facies classification for fluvial sediments

Table 2 *Facies classification. From Miall (1978b).*


Facies code	Facies	Sedimentary Structures	Interpretation
Gms	massive, matrix supported gravel	grading	debris flow deposits
Gm	massive or crudely bedded gravel	horizontal bedding, imbrication	longitudinal bars, lag deposits, sieve deposits
Gt	gravel, stratified	trough cross beds	minor channel fills
Gp	gravel, stratified	planar cross beds	longitudinal bars, deltaic growths from older bar remnants
St	sand, medium to very coarse, may be pebbly	solitary or grouped trough cross beds	dunes (lower flow regime)
Sp	sand, medium to very coarse, may be pebbly	solitary or grouped planar cross beds	linguoid, transverse bars, sand waves (lower flow regime)
Sr	sand, very fine to coarse	ripple cross lamination	ripples (lower flow regime)
Sh	sand, very fine to very coarse may be pebbly	horizontal lamination parting or streaming lineation	planar bed flow (upper flow regime)
Sl	sand, very fine to very coarse may be pebbly	low angle (<10°) cross beds	scour fills, washed-out dunes, antidunes
Se	erosional scours with intraclasts	crude cross bedding	scour fills
Ss	sand, fine to very coarse, may be pebbly	broad, shallow scours	scour fills
Fl	sand, silt, mud deposits	fine lamination, very small ripples	overbank or waning flood
Fsc	silt, mud	laminated to massive	backswamp deposit
Fcf	mud	massive, with freshwater molluscs	backswamp pond deposits
Fm	mud, silt	massive, desiccation cracks	overbank or drape deposits
C	coal, carbonaceous mud	plant, mud films	swamp deposits
P	carbonate	pedogenic features	paleosol

Facies interpretation is often facilitated by considering the vertical facies succession. Where there is a conformable vertical succession of facies, with no major breaks, the facies are the products of environments which were originally laterally adjacent.

This concept has been appreciated since Johannes Walther expounded his Law of Facies in 1894. **The vertical succession of facies is produced by the lateral migration of one environment over another (e.g. the progradation of a delta or tidal flat). Where there are breaks in the succession, seen as sharp or erosional contacts between facies, then the facies succession need not reflect laterally adjacent environments but could well be the products of widely separated environments.**



Examples of fining-upward and coarsening-upward facies successions




Boggs (2001), p.260

Fluvial example of a fining-upward sequence

(a)	<i>Main facts</i>	<i>Processes</i>	<i>Environment</i>
	Coarse, red, structureless siltstone. No body fossils. Sparse concretions of calcium carbonate. Invertebrate burrows in lower part. Fine grained red sandstone. Ripple drift cross laminated. Invertebrate burrows. Fine to medium grained red sandstone. Trough cross bedded with unidirectional palaeocurrents. Siltstone clasts on set bases. Convolute bedding in some sets. Upwards reduction in set size. Sharp surface with slight relief. Scoured sole marks. Concentration of siltstone clasts above surface.	Deposition of silt from suspension. Could be from air or water. Post-depositional burrowing and growth of concretions. Rapid deposition of sand during bedload transport as current ripples. Post-depositional colonisation by animals. Net accumulation of sand during migration of dunes. Post-depositional liquefaction suggests rapid deposition. Erosion taking place nearby. Erosion by fluid scour. Winnowing of finer sediment to leave a lag of intraformational clasts.	Continental, overbank interchannel setting accumulating sediment by vertical accretion. Concretions grow in caliche soil profile. Channel margin/levee? Laterally migrating channel with deposition on one side; probably a point bar of a meandering stream.

Deltaic example of a coarsening-upward sequence

(b)	Main facts		Environment
	Cross bedded sandstone. Sharp contact with relief and siltstone clasts. Siltstone with parallel lamination and graded sand beds. Siltstone. Mudstone with marine fossils. Coarsening upwards unit. Mudstone. Siltstone with current ripples, burrows on top. Cross laminated sandstone with silty partings. Silty mudstone with siltstone laminae. Gradationally striped silty mudstone. Mudstone with marine fossils.	Dune migration; strong currents. Erosion and winnowing. Deposition from suspension, with episodic decelerating flows. Deposition from suspension. Deposition from suspension; high salinity. Increase in energy; shallowing? Deposition from suspension. Mixture of deposition from suspension and from bedload transport by weak currents. Bedload transport as ripples with quiet interludes. Deposition from siltstone with fluctuating supply. Deposition from suspension.	prograding delta front Fluvial distributary channel. Mouth bar. Delta slope. Minor readvance of delta? Abandonment of mouth bar. prograding delta front Proximal mouth bar. Delta slope. Offshore; pro-delta.

Progressive facies successions are commonly terminated by **bounding discontinuities** of some type which generally characterized by abrupt changes in lithology across discontinuities or abrupt changes in facies; these might be erosion surfaces, or surfaces of non-deposition.



Simplified classification of ancient depositional environments

Primary Depositional Setting	Major Environment	Subenvironment
Continental	*Fluvial	{ *Alluvial fan *Braided stream *Meandering stream
	*Desert	
	Lacustrine	
	*Glacial	
Marginal-marine	*Deltaic	{ *Delta plain *Delta front *Prodelta
	*Beach/barrier island	
	*Estuarine/lagoonal	
	Tidal flat	
Marine	Neritic	{ Continental shelf **Organic reef
	Oceanic	{ Continental slope Deep-ocean floor

*Dominantly siliciclastic deposition

**Dominantly carbonate deposition

Boggs (2006), p.243

Environments not marked by an asterisk(s) may be sites of siliciclastic, carbonate, evaporite, or mixed sediment deposition depending upon conditions.

Table 8–1 Criteria for Recognition of Ancient Sedimentary Environments

Criteria based on primary depositional properties

Mainly physical properties.

Geometry of facies units—useful only if very distinctive, e.g., ribbon shape of channels; lobate shape of deltaic deposits.

Gross lithology and mineralogy of strata—a very general environmental indicator; e.g., fossiliferous limestone suggests shallow marine shelf settings; coal indicates swampy environments; the mineral glauconite suggests marine conditions.

Facies associations (stratigraphic successions)—for example, fining-upward successions are characteristic of meandering-stream deposits; regressive shelf environments (shoreline advancing seaward with time) produce coarsening-upward successions.

Sedimentary structures.

Nondirectional structures—not unique environmental indicators but suggest depositional process; e.g., ripples indicate current flow, graded bedding indicates settling of grains from suspension, mudcracks indicate subaerial exposure.

Directional structures (paleocurrent indicators)—paleocurrent patterns may have environmental significance; e.g., bimodal patterns suggest tidal influence; unimodal patterns of high variability suggest meandering-stream environments.

Sedimentary textures—grain-size data of limited usefulness; grain shape measured by Fourier analysis may be significant; grain orientation (e.g., imbrication) a useful paleocurrent indicator.

Boggs (2001), p.261

Mainly chemical properties

Major-element composition—very limited usefulness.

Trace-element composition—some application in paleosalinity interpretation; e.g., boron more abundant in marine shales than in freshwater shales.

Isotope ratios—carbon and oxygen isotopes may be used to interpret marine vs. nonmarine conditions; oxygen isotopes a possible ocean paleotemperature indicator.

Mainly biologic properties.

Kinds of fossils and their ecologic characteristics—very useful indicators of salinity, temperature, depths, energy, and turbidity of ancient oceans; also an indicator of substrate type (rock, sand, mud).

Types of trace fossils—water depth indicators.

Criteria based on derived sediment properties

Properties measured or interpreted from instrumental well logs.


Properties such as rock resistivity, velocity of sound transmission, and natural radioactivity can be measured in well bores and used, for example, to interpret coarsening- and fining-upward successions in subsurface strata.

Characteristics interpreted from seismic reflection records.

Seismic reflection characteristics identified from seismic records indicate features such as inclined bedding, truncations, and pinchouts that have environmental significance.

8. Continental Environments


In the **continental environment**, most sediment transport is accomplished by flows that move in response to the gravity field from positions of greater to lesser potential energy except wind-blown sediment. Where the flow paths of wind flows are determined by the local barometric gradient rather than the earth's gravity field.




沈積環境示意圖。陸棚水深200m以內，陸坡深200m至洋盆。注意陸坡坡度只有4-5度，本圖因有垂直放大，使陸坡坡度似達80度。(鄧屬予，1997)

8.1 Alluvial Fans (沖積扇)


DEBRIS-FLOW-DOMINATED FAN



BRAIDED FLUVIAL FAN



LOW-SINUOSITY/MEANDERING FLUVIAL FAN




Stanistreet & McCarthy (1993)





Alluvial fans are deposits with gross shapes approximating a segment of a cone and exhibiting a convex-up transverse profile.

Figure 8.1

Aerial view of a debris-flow-dominated alluvial fan at the mouth of a canyon in the steep east wall of Death Valley, California. The highway gives the scale. [Photograph by John Shelton.]



Boggs (2006), p.246




Debris-flow versus stream-flow-dominated alluvial fans

Boggs (2006), p.247


Figure 8.2

Schematic diagrams illustrating the depositional features of (A) debris-flow and (B) stream-flow-dominated alluvial fans adjacent to active normal faults. [Modified from Blair and McPherson, 1994, Alluvial fans and their natural distinction from rivers based on morphology, hydraulic process, sedimentary processes, and facies assemblages: Jour. Sedimentary Research, v. A34, Fig. 1, p. 455, reproduced by permission of the Society for Sedimentary Geology.]

Typical surface features and cross-sectional profiles of alluvial fans



8.2 River Systems



Galloway & Hobday (1996)







Figure 8.6
Braid bars in a braided lower reach of the Kongakut River, Arctic National Wildlife Refuge, northeastern Alaska.


Braided Rivers

Boggs (2006), p.251



Boggs (2006), p.253


Structure of bars and vertical sequences in **braided rivers**



Galloway & Hobday (1983)

Figure 8.7

Landsat photograph of the Bramaputra River immediately north of its confluence with the Ganges, showing anastomosing, single-channel, and braided channel patterns. [From Bridge, J. S., 1993, The interaction between channel geometry, water flow, sediment transport and deposition in braided rivers, in Best, J. L., and C. S. Bristow (eds.), Braided rivers, Geological Society London Special Publication No. 75, Fig. 4, p. 21, reproduced by permission.]



Typical facies and vertical profiles of braided rivers:

(A) shallow, gravel-bed braided river;

(B) sheetflood distal braided river

Fluvial architecture of braided-river deposits

Boggs (2006), p.256




Figure 8.11

Schematic representation of the fluvial architecture of braided-river deposits. [After Walker, R. G., and D. J. Cant, 1984, Sandy fluvial systems, in R. G. Walker (ed.), *Facies models*: Geoscience Canada Reprint Ser. 1, Fig. 9, p. 77, reprinted by permission of Geological Association of Canada.]





Gravelly braided-river deposits

Sharp erosional contact

Sandy braided-river deposits

頭料山層(草屯)

Morphological elements of a meandering-river system



Note: A thalweg is a line connecting the deepest points along a stream channel; it is commonly the line of maximum current velocity.





Figure 8.8
Helical flow in a meander bend, leading to lateral accretion of cross-bedded, fining-upward deposits. [From Leeder, M., 1999, *Sedimentology and sedimentary basins*, Fig. B17.1, p. 313. Reproduced by permission of Blackwell Science Ltd.]

Helical flow in a meander bend



Fluvial architecture of meandering-river deposits

Typical lithofacies and vertical profiles (A) Sandy meandering river; (B) fine-grained meandering river





Figure 8.12
Schematic representation of the fluvial architecture of meandering-river deposits. [After Walker, R. G., and D. J. Cant, 1984, *Sandy fluvial systems*, in R. G. Walker (ed.), *Facies models: Geoscience Canada Reprint Ser. 1*, Fig. 9, p. 77, reprinted by permission of Geological Association of Canada.]

Miall (1996)

Overbank Deposits



The majority of Sabunchi sediment (Pliocene, Russia) represents deposition adjacent to channels and channel belts in **flood-plain** facies. Sand bodies are thin and reduce thickness, connectivity, and grain size along strike away from channels. Siltstones and muddy siltstones are common and are continuous across the outcrops.


Paleosols



Despite rapid accumulation rates, pedogenesis is common in the Sabunchi. Pedogenic features are most pronounced within the distal overbank facies, but occur in other environments as well.

Reddish and red-brown horizons typically contain several paleosol indicators including glaes, horizonization, precipitated minerals, and rooting. Most of the soil features are relatively immature, reflecting rapid sediment accumulation and burial.

Paleosols



Despite rapid accumulation rates, pedogenesis is common in the Sabunchi. These inceptisols show rooting and weak horizonization associated with mineral precipitation.

Overbank Deposits



These mud and silt prone intervals contain interbedded thin sands. Mottling, coloration, and rooting suggest common and cyclic paleosol formation. ²⁶

Multistory Channels



Numerous, thick-bedded, multi-story fluvial packages crop out within the succession, commonly above sequence boundaries. These range in thickness from 6 to 40 meters of medium- to fine-grained sandstones (occasionally coarse grained).

Multistory Channels



Multistory, sand-filled channels was cut by shale-filled channels (dash line indicates channel bases).

Channel Sands



The channel sandstones themselves comprise trough-crossbedded, planar bedded, and low-angle inclined facies. These facies may change across or within an individual channel. These high energy elements contrast with the climbing-ripple dominated sands bodies of the Sabunchi or Kirmaky Suites.

Channel Sands

Individual troughs can be as large as 1 m in height and reflect a moderate spread of paleocurrent azimuths.



In both these cases (and others), trough lamina sets are cut by root or burrowing traces.